

AD-A122 107

SALIVARY AND PLASMA TESTOSTERONE AND CORTISOL DURING  
MODERATELY HEAVY EXERCISE(U) NAVAL HEALTH RESEARCH  
CENTER SAN DIEGO CA R R VICKERS ET AL. APR 82  
NAVHLTHRSCHC-82-5

1/1

UNCLASSIFIED

F/G 6/5


NL



END

FILED

DTIC



MICROCOPY RESOLUTION TEST CHART  
NATIONAL BUREAU OF STANDARDS-1963-A

AD A 122107

SALIVARY AND PLASMA TESTOSTERONE AND CORTISOL  
DURING MODERATELY HEAVY EXERCISE

Ross R. Vickers, Jr., Ph.D.<sup>1</sup>

James A. Hodgdon, Ph.D.<sup>1</sup>

Brad L. Bennett, M.S.<sup>2</sup>

Russell E. Poland, Ph.D.<sup>3</sup>

Robert T. Rubin, M.D., Ph.D.<sup>3</sup>

Report No. 82-5, supported by Naval Medical Research and Development Command, Bethesda, Maryland, Department of the Navy, under research Work Unit M0096.PN001-1035, under Office of Naval Research Contract N00014-81-K-0561; and by NIMH Research Scientist Development Award MH47363 (to R.T.R.). The views presented in this paper are those of the authors. No endorsement by the Department of the Navy has been given or should be inferred.

<sup>1</sup>Environmental Physiology Department, Naval Health Research Center, P.O. Box 85122, San Diego, CA 92138

<sup>2</sup>Navy Submarine Medical Research Laboratory, New London, CT.

<sup>3</sup>Department of Psychiatry, Harbor-U.C.L.A. Medical Center, Torrance, CA.

The authors thank Cheryl Spinweber and Earl A. Edwards for their helpful suggestions and comments on an earlier draft of this paper.

have been reported. Studies have shown the relationship of steroid hormones. High correlations between salivary and urinary levels of cortisol, corticosterone, and 17-ketotestosterone have been reported under various conditions of activity, including periods of sleep, indicating some of the steroid hormones. However, the relationship between salivary and urinary levels of these hormones is not always consistent. The present study was designed to investigate the relationship of salivary and plasma measures of cortisol and corticosterone during exercise.

Seven subjects received a baseline test at 7 percent of their maximal oxygen consumption capacity. Paired blood and saliva samples were obtained at the beginning of the test, at the beginning of the run, at 15 minutes into the run, and at the end of the run. Blood concentrations and cortisol values were determined for each sample by radioimmunoassay. Plasma corticosterone concentrations were determined by the same technique. The relationship between the salivary and plasma concentrations of cortisol and corticosterone was determined at the end of the run during the study.

Correlations between salivary and plasma measures were calculated (Spearman's  $r = .45$  to  $r = .55$ ). Significant correlations were found between salivary and plasma measures of cortisol and corticosterone in either plasma or saliva due to large interindividual differences in concentrations. Significant correlations were found during exercise for the salivary measures.

Salivary measures of cortisol and corticosterone were found to be reliable measures of the relationship between salivary and plasma measures of these hormones. The relationship between salivary and plasma measures of these hormones was found to be similar to that reported in previous studies. The relationship between salivary and plasma measures of these hormones was found to be similar to that reported in previous studies. The relationship between salivary and plasma measures of these hormones was found to be similar to that reported in previous studies. The relationship between salivary and plasma measures of these hormones was found to be similar to that reported in previous studies.

## Introduction

The study of steroid hormones is of interest to exercise physiologists because these hormones influence metabolic substrate utilization and fluid and electrolyte balance. However, changes in levels of circulating steroid hormones during exercise have been inconsistent across studies (1-4). Resolution of these inconsistencies might be facilitated by more frequent sampling of the hormones to accurately quantify the temporal profile of hormone changes during exercise. Techniques that facilitate frequent sampling therefore may contribute to improved understanding of the physiological processes involved in exercise.

Standard blood collection procedures can be difficult to employ when frequently repeated measures are needed. Drawing blood for these measures can be troublesome due to problems arising from the need for repeated venipunctures. Catheterization provides an alternative, but is impractical in many settings. Venipuncture by either method may disturb the hormones to be measured (5).

Salivary measures provide a potentially sensitive alternative means of monitoring steroid hormones. Close parallels between salivary concentrations and plasma or serum concentrations have been demonstrated for cortisol(6-9), testosterone (10-13), 17-hydroxyprogesterone (14-16), and aldosterone (17). Salivary measures have the advantages of low risk, minimal stress, and ease of repetition during short time periods. However, this work has been done in clinical settings under resting conditions, and the findings may not generalize to conditions of interest to exercise physiologists. Thus, the present study was undertaken to examine the impact of a 30-minute period of moderately heavy exercise on the association between plasma and salivary measures of cortisol and testosterone.

## Method

### Sample

Fifteen male military and civilian personnel (12 white, three black) working for the U.S. Navy in San Diego, California, participated after giving informed consent and passing a medical screening examination. Additional descriptive data are given in Table 1. (See page 4.)

### Exercise Protocol

The research design required participants to complete a 30-minute run at 75% of maximum oxygen consumption capacity ( $\text{VO}_2 \text{ max}$ ). Participants reported to the laboratory between 1300 and 1330 for  $\text{VO}_2 \text{ max}$  assessment, which was measured in a single graded exercise session on a treadmill. Participants walked at 3.0 miles per hour (mph) at 0% grade for four minutes to warm up. The speed then was increased 0.5 mph each minute until a comfortable pace between 7 and 8 mph was reached. The grade then was increased 3% per minute until the participant voluntarily stopped running. Spirometry values during the run were determined by an online, open-circuit spirometry system<sup>1</sup> which recorded values on a magnetic tape and printed out 15-second averages.  $\text{VO}_2 \text{ max}$  was reached when  $\text{VO}_2$  did not increase in the one minute average during the test. The measured  $\text{VO}_2 \text{ max}$  indicated that participants were above average in fitness (Table 1).

---

<sup>1</sup>The spirometry system consisted of an Applied Electrochemistry  $\text{O}_2$  analyzer (Model S3-A), A Beckman  $\text{CO}_2$  analyzer Model (LB-2), a KL Engineering pneumoscan spirometer (Model S-300), a Yellow Springs air temperature thermometer (Model 43T-A) and an Bwald Koegel one-way valve. All interfaced with a Hewlett-Packard Model 9825A programmable desk-top calculator.

TABLE 1  
PARTICIPANT DESCRIPTION

	MEAN <sup>a</sup>	STANDARD DEVIATION
Age (years)	28.1	4.5
Height (cm)	177.2	8.5
Weight (kg)	78.6	10.4
Sum of skinfolds <sup>b</sup>	29.2	10.6
VO <sub>2</sub> max (l/min)	4.48	0.68
VO <sub>2</sub> (ml/kg min)	58.4	4.4
<i>PHYSIOLOGICAL RESPONSES TO THE 30-MINUTE TREADMILL RUN</i>		
Heart rate (bpm)	165.5	11.3
Ventilation (l/min STPD)	92.8	18.8
VO <sub>2</sub> (l/min STPD)	3.49	0.50
VO <sub>2</sub> (ml/kg min STPD)	45.6	2.9
VO <sub>2</sub> (% VO <sub>2</sub> max)	77.5	4.2

<sup>a</sup>N = 15.

<sup>b</sup>Computed from measures of biceps, triceps, suprailiac, and subscapular skinfold.

The day following VO<sub>2</sub> max determination, the participants returned to the laboratory at the same time of day for the 30-minute run. Each participant ran at a speed calculated to represent 75% of his VO<sub>2</sub> max. On the average, actual VO<sub>2</sub> during the run was quite close to 75% (Table 1). This work level was expected to produce testosterone and cortisol changes in moderately conditioned individuals (1,2). Prior to beginning the run, the participants provided blood and saliva samples (see below) and then warmed up at six mph for five minutes. They then ran for 15 minutes, gave another blood and saliva sample, and completed the last 15 minutes of the run. At the end of the run, final blood and saliva samples were collected.

#### Blood and Saliva Collection and Assay Procedures

Venous blood was drawn from a superficial vein of the antecubital fossa into a 15-ml vacutainer collection tube. Samples were centrifuged immediately and the plasma separated from the red blood cells and frozen. Three-minute samples of whole saliva were collected by having the participants expectorate directly into 35-ml containers. Salivation was stimulated by chewing a 1.0 gm bolus of paraffin. Samples were frozen immediately after collection. The time required to collect both samples was three to four minutes, as blood was drawn during the saliva collection. In addition to the three samples taken during the 30-minute run, a baseline measure was taken the day before the run during the participants' initial laboratory visit prior to the VO<sub>2</sub> max determination.

To determine flow rate, saliva volume was estimated as follows: The vacutainer was first weighed empty. Empty weight was subtracted from the combined weight of the sample and the vacutainer to determine net sample weight, which then was translated to a volume estimate by assuming that the specific gravity of the saliva was 1.000. This assumption should have introduced an error of 1 percent at most into the volume estimates (18, pp. 22-23). Such an error is very small compared to the observed differences in estimated sample volume. Flow rate was finally determined as milliliters per minute by dividing the estimated volume by three.

The samples were stored frozen until assayed. All samples were measured in a single assay. Cortisol radioimmunoassay (RIA) employed the technique of Foster and Dunn (19) with intra-assay coefficients of variation of 17.6%, 5.2%, and 14.0% for low, medium, and high concentration pools, respectively. The testosterone RIA used the technique of Odell, et al. (20) with intra-assay coefficients of variation of 17.2%, 8.8%, and 2.0% for low, medium, and high concentration control pools, respectively. The RIAs measured total blood and total salivary hormone concentrations.

#### Analysis Procedures

Pearson product-moment correlations were used to describe the association between plasma and salivary concentrations for cortisol and testosterone. Additional analyses ensured that these product-moment correlations accurately summarized the data. The Kolmogoroff-Smirnov test (21) indicated that the score distributions met the assumption of normality. Nonparametric correlation (21) and jackknife statistics (22) showed that the parametric statistics yielded reasonable estimates of magnitude of association and significance, even if the normality assumptions had been relaxed. A multivariate analysis of variance for repeated measures was used to determine whether mean hormone levels changed significantly during the 30-minute run. All analyses except the jackknife statistics were performed with the Statistical Package for the Social Sciences (SPSS) (23). The jackknife statistics were computed on a hand calculator from correlations produced by the SPSS program.

#### Results

Cortisol levels did not change significantly during the study, but testosterone levels did (Table 2). Examination of values for individual participants showed that the nonsignificant cortisol trends represented the effect of averaging over people with very different response patterns. In the case of testosterone, there was a general correspondence in the trends for mean plasma and mean saliva hormone concentrations.

TABLE 2  
PLASMA AND SALIVA CONCENTRATIONS OF CORTISOL AND TESTOSTERONE:  
MEAN LEVELS AND INTERCORRELATIONS

			BASELINE	PRE-EX	15 MINUTES	30 MINUTES	AVERAGE	F VALUE	SIGNIFICANCE
CORTISOL	Plasma	Mean <sup>a</sup>	84.40	94.83	91.87	95.27	91.62	2.17	.145
		S. D.	22.72	32.44	29.77	33.81	29.56		
	Saliva	Mean <sup>a</sup>	1.89	2.31	2.12	2.47	2.20	1.84	.194
		S. D.	.88	1.54	1.06	1.60	1.30		
	Plasma/Saliva Correlation		.82***	.87***	.86***	.95***	.88***		
TESTOSTERONE	Plasma	Mean <sup>a</sup>	5.37	5.54	6.58†	6.73†	6.06	5.40	.014
		S. D.	1.74	1.58	2.60	2.21	2.11		
	Saliva	Mean <sup>a</sup>	.079	.085	.105	.115†	.096	4.19	.030
		S. D.	.033	.019	.030	.036	.033		
	Plasma/Saliva Correlation		.65**	.24	.48*	.37	.51*		

<sup>a</sup>All concentrations reported in ng/ml

\* $p < .05$ , one-tailed

\*\* $p < .01$ , one-tailed

\*\*\* $p < .001$ , one-tailed

†Significantly different ( $p < .05$ ) from baseline and pre-exercise values



Paired plasma and saliva cortisol concentrations were highly correlated ( $r_{\text{c}} \geq .82$ , Table 2). However, plasma:saliv. testosterone correlations were consistently lower ( $r_{\text{c}} \leq .65$ , Table 2).

Salivary cortisol concentrations ranged between 2.14 and 2.36 percent of the corresponding plasma concentrations. Corresponding values for testosterone were 1.48 to 1.79 percent. In both cases, there was a nonsignificant trend toward lower ratios during the later sample collections. Salivary hormone concentrations were not correlated with saliva flow rate at any point.

#### Discussion

Moderately heavy exercise did not affect the relationship between plasma and salivary cortisol. The correlations were consistently high. Mean levels of cortisol did not vary, possibly due to the only moderate exercise intensity (3). Whether this will hold true during heavier or more prolonged exercise is yet to be determined.

The testosterone findings were equivocal. Paired saliva and plasma values were not highly correlated. Changes in mean testosterone levels during exercise were significant and similar in pattern for both plasma and saliva. The low correlation between the paired samples is at odds with previous findings in resting subjects (13), but even the baseline correlation in this study was only  $r = .65$ . One aspect of our data that might account for this discrepancy was a relatively restricted range of testosterone values compared to those in other studies. Wang, et al. (13) reported a high correlation ( $r = .94$ ) and presented a scatterplot for paired saliva and plasma testosterone values. The scatterplot suggests that elimination of a few extreme samples would have produced results quite similar to ours. Within a restricted range of values, the well-documented individual differences in the ratio of free to total testosterone (24-26) could have had more influence on the saliva-plasma associations. This issue may be important for salivary steroid research in general because similar distributional factors appear to affect some of the reported associations for cortisol (16), 17-hydroxyprogesterone (16), and aldosterone (17). Previous reports of high salivary-plasma or salivary-serum correlations therefore may have depended on the inclusion of samples from persons with endocrine problems or who received drugs that altered steroid levels.

The present findings do not rule out the possibility that salivary testosterone might have been highly correlated with the free plasma testosterone fraction in our subjects. It is commonly held that salivary steroids reflect free plasma steroids more than they do total plasma steroids (6-8,10-13). Unfortunately, this belief is based largely on indirect evidence (e.g., absence of steroid binding globulins in saliva, low salivary steroid concentrations, and similar half-lives of free plasma and salivary steroids). Katz and Shannon (9) have shown that salivary corticosteroids were more highly correlated with free plasma corticosteroids ( $r = +.83$ ) than with total plasma corticosteroids ( $r = +.75$ ), but this difference is too small for final conclusions, even with circumstantial support from other lines of investigation. The fact that total plasma values and salivary values can be relatively independent at times, as in the case of our testosterone data, makes it critical to examine this issue in more detail. Unfortunately, the importance of the free vs. total blood hormone distinction was not anticipated at the outset of this project because of the consistently high correlations reported in other research; therefore, free hormone measures are not available in our data. Despite the expense and difficulty of obtaining both free and total hormone values, further investigation of the differences between the two is essential. This is particularly true if salivary measures are to be employed in studies involving normal populations.

The fact that testosterone levels increased with exercise adds additional data to support the position that testosterone concentrations increase during exercise; previous evidence on this issue has been mixed (2-4).

The main conclusion from our study is that salivary cortisol can be substituted for plasma cortisol in research involving moderately heavy exercise. The equivocal testosterone findings point to a need for direct investigation of

the relationship between salivary steroids and both total and free plasma steroids; this need applies to other steroid hormones as well.

#### References

1. Shephard, R.J. & Sidney, K.H. Effects of physical exercise on plasma growth hormone and cortisol levels in human subjects. In J.H. Wilmore & J.F. Keogh (Eds.), Exercise and Sport Science Reviews, 1975, Academic Press, pp. 1-30.
2. Sutton, J.R., Coleman, M.J. & Casey, J.H. Testosterone production rate during exercise. In Landry, F. & Orban, W.A.R. (Eds.), Third International Symposium on Biochemistry of Exercise, 1978, Symposia Specialists, Inc.
3. Aamundsen, L.R. & Balke, B. Glucocorticoid responses to acute and chronic physical exercise. Medicine and Science in Sports, 1973, 5, 59.
4. Kuoppasalmi, K., Naveri, H., Kosunen, K., Harkonen, M., & Adlercreutz, H. Plasma steroid levels in muscular exercise. In Poortmans, J. & Niset, G. (Eds.), Biochemistry of Exercise IV-B, 1981, University Park Press, Baltimore, MD, pp. 149-160.
5. Rose, R.M. & Hurst, M.W. Plasma cortisol and growth hormone responses to intravenous catheterization. Journal of Human Stress, 1975, 1, 22-36.
6. Ueda, T., Hiramatsu, R., Iwaoka, T., Shimada, T., Miura, F., & Sato, T. Use of saliva for monitoring unbound free cortisol levels in serum. Clinica Chimica Acta, 1981, 110, 245-253.
7. Streat, L.P. Variation in salivary levels of 17-hydroxycorticosteroids as an index of adrenocortical activity and circulating corticosteroids. The New York State Dental Journal, 1960, 26, 368-370.
8. Walker, R.F., Riad-Rahmy, D., & Read, G.F. Adrenal status assessed by direct radioimmunoassay of cortisol in whole saliva or parotid saliva. Clinical Chemistry, 1978, 24, 1460-1463.
9. Katz, F.H. & Shannon, I.L. Parotid fluid cortisol and cortisone. Journal of Clinical Investigation, 1969, 48, 838-855.
10. Landman, A.D., Sanford, L.M., Howland, B.E., Dawes, C., & Pritchard, E.T. Testosterone in human saliva. Experientia, 1976, 32, 940-941.
11. Luisi, M., Bernini, G.P., Del Genovese, A., Birindelli, R., Barletta, D., Gasperi, M., & Franchi, F. Radioimmunoassay for "free" testosterone in human saliva. The Journal of Steroid Biochemistry, 1980, 12, 513-516.
12. Smith, R.G., Besch, P.K., Dill, G., & Buttram, Jr., V.C. Saliva as a matrix for measuring free androgens: Comparison with serum androgens in polycystic ovarian disease. Fertility and Sterility, 1979, 31, 512-517.
13. Wang, C., Plymate, S., Nieschlag, E., & Paulsen, A. Salivary testosterone in men: Further evidence of a direct correlation with free serum testosterone. Journal of Clinical Endocrinology and Metabolism, 1981, 53, 1021-1024.
14. Walker, R.F., Read, G.F., Hughes, I.A., & Riad-Rahmy, D. Radioimmunoassay of 17 $\alpha$ -hydroxyprogesterone in saliva, parotid fluid, and plasma of congenital adrenal hyperplasia patients. Clinical Chemistry, 1979, 25, 542-545.
15. Walker, R.F., Read, G.F., & Riad-Rahmy, D. Radioimmunoassay of progesterone in saliva: Application to the assessment of ovarian function. Clinical Chemistry, 1979, 25, 2030-2033.
16. Price, D.A., Astin, M.P., Chard, C.R., & Addison, G.M. Assay of hydroxyprogesterone in saliva. Lancet, 18 Aug 1979, 368-369.
17. McVie, R., Levine, L.S., & New, M.I. The biologic significance of aldosterone concentration in saliva. Pediatric Research, 1979, 13, 755-759.

18. Kerr, A.C. The Physiological Regulation of Salivary Secretions in Man. New York: Pergamon Press, 1961.
19. Foster, L.B. & Damm, R.T. Single antibody technique for radioimmunoassay of cortisol in unextracted serum or plasma. Clinical Chemistry, 1974, 20, 365-368.
20. Odell, W.D., Swerdloff, R.S., Bain, J., Wollesen, F., & Grover, P.K. The effect of sexual maturation on testicular response to LH stimulation of testosterone secretion in the intact rat. Endocrinology, 1974, 95, 1380-1384.
21. Siegel, S. Nonparametric Statistics for the Behavioral Sciences. New York: McGraw-Hill, 1956.
22. Duncan, G.T. & Layard, M.W.J. A Monte-Carlo study of asymptotically robust tests for correlation coefficients. Biometrika, 1973, 60, 551-558.
23. Nie, N.H., Hull, C.H., Jenkins, J.G., Steinbrenner, K., & Bent, D.H. Statistical Package for the Social Sciences. New York: McGraw-Hill Book Company, 1975.
24. Vermeulen, A., Stoica, T., & Verdonck, L. The apparent free testosterone concentration, an index of androgenicity. Journal of Clinical Endocrinology, 1971, 33, 759-767.
25. Moll, Jr., G.W., Rosenfield, R.L., & Helke, J.H. Estradiol-testosterone binding interactions and free plasma estradiol under physiological conditions. Journal of Clinical Endocrinology and Metabolism, 1981, 52, 868-874.
26. Chopra, I.J. & Tulchinsky, D. Status of estrogen-androgen balance in hyperthyroid men with Graves' disease. Journal of Clinical Endocrinology and Metabolism, 1974, 38, 269-277.

Unclassified

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER 82-5	2. GOVT ACCESSION NO. AD-A122107	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle)  Salivary and Plasma Testosterone and Cortisol During Moderately Heavy Exercise		5. TYPE OF REPORT & PERIOD COVERED  Interim
7. AUTHOR(s)  Ross R. Vickers, Jr., James A. Hodgdon, Brad L. Bennett, Russell E. Poland, R. T. Rubin		6. PERFORMING ORG. REPORT NUMBER
9. PERFORMING ORGANIZATION NAME AND ADDRESS  Naval Health Research Center P. O. Box 85122 San Diego, CA 92138		8. CONTRACT OR GRANT NUMBER(s)
11. CONTROLLING OFFICE NAME AND ADDRESS  Naval Medical Research & Development Command Bethesda, MD 20014		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS  M0096.PN001-1035
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)  Bureau of Medicine & Surgery Department of the Navy Washington, D.C. 20372		12. REPORT DATE April 1982
		13. NUMBER OF PAGES 9
		15. SECURITY CLASS. (of this report)  Unclassified
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report)  Approved for public release; distribution unlimited.		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)  Approved for public release; distribution unlimited.		
18. SUPPLEMENTARY NOTES		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number)  Cortisol                      Saliva Exercise                      Testosterone Plasma		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number)  Saliva may provide a useful alternative to blood for measuring steroid hormones. Total plasma and salivary concentrations of cortisol and testosterone were compared in samples taken twice at rest and twice during exercise to determine whether physical activity level affects the relationship between the two. Correlations were consistently high ( $r > .82$ ) for cortisol, but relatively low for testosterone ( $r < .66$ ). Exercise did not affect either correlation. Salivary cortisol is a reasonable alternative to plasma cortisol		

DD FORM 1 JAN 73 1473

EDITION OF 1 NOV 65 IS OBSOLETE  
S/N 0102-LF-014-6601

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

Unclassified

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

20. Abstract (continued)

even during exercise. The testosterone results were equivocal as salivary testosterone could be highly correlated with free plasma testosterone despite the low correlation to total plasma testosterone. Closer examination of the free/total plasma hormone distinction was not possible in the present study, but should be an important focus for further research on salivary steroids.

A

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

**END**

**FILMED**

**1-83**

**DTIC**